
# **USMLE Step 2CK/3 Rapid Review**

# Written by: Owais Mohammad Ahmad MD

#### **Gyne**

- 1. What is the tx of Endometrial hyperplasia?
  - Progestin therapy (progestin IUD)
- 2. What marker detects that ovarian reserves are aging quickly?
  - Low anti mullerian hormone
  - Please note, anti mullerian hormone produced by the granulosa cells inhibits the
    recruitment and premature growth of ovarian follicles. Thereby playing a significant
    role in the regulation of folliculogenesis (the development of ovarian follicles). By
    preventing the over-activation of follicles, AMH helps maintain a reserve of follicles
    for future ovulation.
- 3. What is the likely diagnosis in an African-American woman at 20 weeks gestation that presents with new-onset hirsutism and acne? Pelvic ultrasound reveals an intrauterine gestation consistent with dates and bilateral 8-cm solid ovarian masses.
  - Luteoma, Please note (A common wrong answer here is Theca lutein cyst) --> Theca lutein cyst is a wrong answer as that is associated w/ a Hyadtiform mole
  - Luteoma is a tumor of the ovary. it is stimulated by bHCG to convert the stromal cells
    to undergo luetinization. Lutenization refers to the biological process during which the
    ovarian follicle transforms into a corpus luteum after ovulation. Luteomas are ovarian
    tumors that can produce and secrete androgens (male hormones such as
    testosterone and androstenedione).



- 4. 27F w/ Anemia and Leiden Factor 5 mutation What is the best contraceptive method for her?
  - Progestin IUD, Please note, estrogen-containing contraceptives (e.g.,COPC, patches, vaginal rings) are contraindicated in this patient as they increase the risk of VTE. Also, Copper IUDs are contra-indicated in this patient as they increase heavy menstrual bleeding. The copper component induces a mild inflammatory reaction within the uterine lining. This inflammatory response increases prostaglandin production, which can enhance uterine contractions and vasodilation, leading to increased blood flow and heavier bleeding.
- 5. What disease can candida cause in the female genitalia?
  - Candida Vaginitis
- 6. What is the MOA Of Raloxifene?
  - Antagonist at breast, uterus; agonist at bone (they are considered partial agonist at the bone)
- 7. What is the MOA of Tamoxifen?

- Antagonist at breast, partial agonist at uterus, bone (they are considered partial agonist at the bone and uterus)
- 8. When does tamoxifen worsen bone loss and when does it improve bone health?
  - It may help <u>postmenopausal women</u> maintain bone density, but better options (e.g., bisphosphonates, denosumab, raloxifene) exist for osteoporosis prevention and treatment.
  - X Tamoxifen can worsen bone loss in <u>premenopausal women</u>, so it is not recommended in younger patients at risk for osteoporosis.
- 9. How many months of unprotected intercourse to diagnose infertility in a woman less than 35 years
  - 12 months of unprotected intercourse
  - Women under 35 generally have:
    - Better ovarian reserve (more and healthier eggs),
    - Higher natural fertility rates per cycle,
    - More time to conceive naturally before age-related decline becomes significant.
  - Thus, waiting 12 months allows a reasonable chance for natural conception without premature interventions
- 10 . How many months of unprotected intercourse to diagnose infertility in a woman above 35 years?
  - 6 months of unprotected intercourse
  - Why 6 months for women 35 or older?
  - Starting around age 35, and especially after 37, fertility declines significantly due to:Decreased ovarian reserve and oocyte quality, Increased rates of anovulation, miscarriage, and chromosomal abnormalities.

The shorter timeframe (6 months) ensures earlier intervention, since:

Delaying evaluation may reduce the chances of successful treatment,

- 11. What is the relationship between TRH and prolactin?
  - TRH stimulates prolactin
- 12. What effect does alcohol have on the reproductive system?
  - Alcohol causes sexual dysfunction as it decreases GnRH which decreases
     Testosterone
- 13. What does AUB PALM COEIN stand for ?
  - Bleeding from the uterus that is irregular in volume, frequency, duration, or timing and occurs in reproductive-age women (i.e., before menopause).
  - Common causes (classified by PALM-COEIN system):
  - PALM (structural): Polyp, Adenomyosis, Leiomyoma (fibroids), Malignancy/hyperplasia
  - COEIN (non-structural): Coagulopathy, Ovulatory dysfunction, Endometrial disorder, latrogenic (anticoagulants), Not yet classified (chronic endometritis)
  - PALM → structural, COEIN → non structural
  - Disorders affecting endometrial hemostasis: Hemostasis is the process your body uses to stop bleeding. In the uterus, this happens every month when the endometrium is shed (your period). Normally, bleeding is controlled by local factors like vasoconstriction, clot formation, and tissue repair. In some women, the endometrial lining doesn't stop bleeding properly, even though everything else (hormones, uterus structure) looks normal.

- 14. What are the indications of endometrial biopsy for women less than 45?
  - AUB + Unopposed estrogen (obesity, anovulation), failed medical management,
     lynch syndrome (herediatry nonpolyposis coloectecal cancer)
- 14. What are the indications of endometrial biopsy for women 35 or above?
  - Atypical glandular cells on Pap test
  - Atypical Glandular Cells (AGC) on a Pap test are an abnormal finding that suggests changes in the glandular (columnar) cells of the cervix or endometrium. These cells are different from the more commonly found squamous cells and are associated with a higher risk of precancerous or cancerous lesions

Endometrial biopsy indications		
Age ≥45  • Abnormal uterine bleeding • Postmenopausal bleeding		
Age <45	Abnormal uterine bleeding PLUS:  • Unopposed estrogen (obesity, anovulation)  • Failed medical management  • Lynch syndrome (hereditary nonpolyposis colorectal cancer)	
Age <u>≥</u> 35	Age ≥35 • Atypical glandular cells on Pap test	

# 15. What is the clinical uses of Leuprolide?

- Uterine fibroids, endometriosis, precocious puberty, prostate cancer, infertility. Pulsatile for pregnancy, continuous for cancer. And infertility
- Fibroids = Fibroids are benign tumors of the uterine smooth muscle that grow in response to estrogen and progesterone. Effect of leuprolide: By lowering estrogen levels.
- Endometriosis: Hormonal dependence: Endometrial tissue outside the uterus (endometriosis) responds to estrogen, leading to proliferation, inflammation, and pain. Effect of leuprolide: Suppressing estrogen:
- 16. Which SERM is used to prevent osteoporosis?
  - Raloxifene → Unlike tamoxifen, raloxifene does not stimulate the endometrium, reducing the risk of endometrial hyperplasia and cancer.
- 17. What happens to the LH:FSH ratio in PCOS? (increase/decrease/no change)> Increase LH to FSH ratio
  - Increase insulin → INcrease GnRH release from the hypothalamus favors LH over FSH, Elevated LH stimulates theca cells in the ovary to produce more androgens.excess androgens produced by the theca cells decrease the production of sex hormone-binding globulin (SHBG), increasing free androgens, which can further disrupt normal feedback controls, perpetuating high LH levels.0

- 18. They might tell you there's a 38-year-old woman who had a tubal ligation two years ago (i.e., there's no way she's pregnant), but on physical exam she has a uterus consistent with 8 weeks gestation. What is the dx?
  - Adenomyosis → A condition in which endometrial tissue exists within and grows into the uterine wall.

# 19. What is the tx of adenomyosis?

- By suppressing ovulation and stabilizing hormonal levels, OCPs reduce the cyclical stimulation of ectopic endometrial tissue within the myometrium.
- Adenomyosis is characterized by ectopic endometrial tissue within the myometrium, which responds to hormonal stimuli and can produce increased prostaglandins.
   Elevated prostaglandin levels contribute to heavy menstrual bleeding (menorrhagia), dysmenorrhea (painful periods), and inflammation. NSAIDs decrease prostaglandin levels.
- 20. What is the NBS in a person presenting w/ testicular cancer?
  - {{c1::Ultrasonography of the scrotum }} If ultrasound reveals a solid lesion, further work-up with serum tumor markers (eg, alpha-fetoprotein, β-hCG) and screening CT scans is typically required Yours = Trans-scrotal biopsy of the right testis Patients with suspected testicular cancer who receive trans-scrotal biopsy are at increased risk for lymphatic spread and recurrence. As such, radical orchiectomy is preferred for both tissue diagnosis and treatment. However, scrotal ultrasound should be performed prior to either invasive procedure in order to determine if the mass is solid or cystic.
- 21. 10M w/ blue colored mass at the superior pole w/ normal cremasteric reflex dx?
  - Torsion of the appendix of the testes.
  - When it twists (torsion), its blood supply is compromised, leading to ischemia and necrosis. The infarcted, necrotic appendix becomes dark blue or black in color.
  - Cremasteric reflex --> stimulation of the illioinguinal nerve causing contraction of the cremaster muscle --> elevation fo the testis
- 23. NBME 11 Block 3 13F (1) tanner stage 2 for breast and pubic hair (2) Never menstruated before. What is NBS ?

40.	A 13-year-old girl is brought to the physician because her mother is concerned that her daughter has never had a menstrual period. She is at the 80th percentile for height and 75th percentile for weight. Breast and public hair development is Tanner stage 2. Examination shows no other abnormalities. Which of the following is the most appropriate next step in management?
0	A) Reexamination in 1 year
0	B) Pelvic examination
0	C) Karyotype analysis
-0	D) Measurement of serum estradiol concentration
0	E) Transvaginal ultrasonography

Correct = Reexamination in 1 year, Wrong answer = Pelvic exam

24. What is the definition of primary Amenorrhea?

- (1) >15F + Sex feature + never menstruated
- (2) >13F + NO sex feature + never menstruated

24. How often to do a pap smear in women who have had total abdominal hysterectomy w/bilateral salpingo-oophorectomy for symptomatic fibroid?

- Never


× 38	3. A 48-year-old Hispanic American woman returns to the office for a routine gynecologic examination. She is generally healthy and says she has felt well. One year ago, she underwent total abdominal hysterectomy with bilateral salpingo-cophorectomy for treatment of symptomatic leicomyomata uteri. She has been taking estraidol since the procedure. She takes no other medications. Revend P aps mears obtained in the past have all been normal. Vital signs are normal and physical examination shows no abnormalities. The patient asks, "Now that I've had a hysterectomy, how often should a Pap smear be done?" Which of the following is the most appropriate recommendation to the patient for having a Pap smear?		
	A) Each year		
	B) Every 2 years		
	C) Every 3 years		
	D) Every 5 years		
	E) Never	NBME 7 B3	
Co	orrect Answer: E.		
of ev ald Pa	Pap smear testing employs a cervical swab to collect cells from the transformation zone of the cervix and is an important screening tool for the detection of cervical carcinoma. The United States Preventive Services Task Force guidelines recommend screening for cervical carcinoma with Pap smear testing every 3 years between ages 21 and 29, and Pap smear screening with co-testing for human papillomavirus (HPV) every 5 years or Pap smear testing alone every 3 years in women between the ages of 30 and 65. Total abdominal hysterectomy involves the surgical removal of the uterus and the cervix. Patients who have undergone this procedure no longer require cervical cancer screening with Pap smears if the hysterectomy was performed for a beingin the particular such as the procedure of the very control to the page of the procedure of the page		
Inc	correct Answers: A, E	3, C, and D.	
wi	Each year (Choice A) or every 2 years (Choice B) are intensified screening regimens that may be appropriate for certain high-risk patients, such as it with a history of previously resected or ablated cervical intraepithelial neoplasia, Iti/O, or maternal exposure to diethystilibestroit. These intensified screening regimens are not required in a patient who has undergone total abdomling hysterectomy for a benign indication.		
Every 3 years (Choice C) is the current recommended screening regimen for women between the ages of 21 and 29, as well as those both of 30 and 65 if the latter group does not undergo co-testing for HPV. This screening regimen is appropriate if a revious Pap smears have be the patients do not have other symptoms or signs of cervical cancer, such as postcottal bleeding. However, this patient has no remaining or and so does not warrant further Pap smear testing.		r group does not undergo co-testing for HPV. This screening regimen is appropriate if previous Pap smears have been normal and e other symptoms or signs of cervical cancer, such as postcoital bleeding. However, this patient has no remaining cervical tissue	

# 25. Which one has elevated PSA?

- Prostate cancer not BPH

Comparison of benign prostatic hyperplasia & prostate cancer			
	ВРН	Prostate cancer	
Risk factors	• Age >50	Age >40, African     American & family history	
Affected part	Central portion     (transitional zone)	Usually peripheral     zone of prostate but can     be anywhere	
Examination	<ul> <li>Symmetrically enlarged &amp; smooth prostate</li> <li>Can have elevated PSA</li> </ul>	Asymmetrically enlarged,     nodules & firm prostate     Markedly elevated PSA	


**BPH** = benign prostatic hyperplasia; **PSA** = prostate-specific antigen.



# <u>Obs:</u>

Ultrasound ass	Itrasound assessment of gestational age		
Ultrasound parameter	Gestational age (weeks)	Accuracy (days)	
Gestational sac diameter	4.5–6	+/- 5–7	
Crown-rump	7–10	+/- 3	
length	11–14	+/- 5	
Biparietal	14–20	+/- 7	
diameter, head circumference,	21–30	+/- 14	
femur length	>30	+/- 21–28	

Uterine abnormalities septate, bicornuate, didelphys "Bi-": means "two" " "Cornu": means "horn." didelphys means double




- 2. Uterine contraction Length?
  - 30 seconds long every 10 minutes
- 3. What is the management of HIV In a newborn?
  - Less than 1000 Vaginal delivery
  - More than 1000 C section
- 4. What is the cause of endometritis?
  - C section: During a C-section, the uterine cavity is breached, providing an opportunity for bacteria from the lower genital tract, skin, or environment to enter the sterile uterine environment. This increases the risk of infection.
- 5. What is the underlying cause of chorioamnionitis?
  - Prolonged ROM: Rupture creates a pathway for bacteria from the lower genital tract (vagina and cervix) to ascend into the sterile intrauterine environment. The longer the membranes remain ruptured, the greater the opportunity for bacterial colonization and infection.
- 6. What are the indications for giving GBS Prophylaxis?
  - Prior infant w/ GBS SEPSIS
  - Rectovaginal Culture positive for GBS in CURRENT pregnancy
  - Unknown status
    - Fever more than 37 C
    - ROM > 18 hours


- Intrapartum fever
- GBS Bacteruria or GBS UTI in current pregnancy (regardless of tx)

Preventing neonatal group B Streptococcus infection	
Antenatal screening	Rectovaginal culture at 36-38 weeks gestation
Indications for intrapartum prophylaxis	<ul> <li>GBS bacteriuria or GBS urinary tract infection in current pregnancy (regardless of treatment)</li> <li>GBS-positive rectovaginal culture in current pregnancy</li> <li>Unknown GBS status PLUS any of the following:         <ul> <li>&lt;37 weeks gestation</li> <li>Intrapartum fever</li> <li>Rupture of membranes for ≥18 hours</li> </ul> </li> <li>Prior infant with early-onset neonatal GBS</li> </ul>
Intrapartum prophylaxis	Intravenous penicillin

If the GBS status is unknown and the person has any one of the following risk factors during labor, intrapartum antibiotic prophylaxis (IAP) (typically intravenous penicillin) is recommended to help prevent neonatal GBS infection:

- 1. Less than 37 weeks gestation Preterm labor increases the risk of passing GBS to the baby. Preterm hence there will be less antibodies for protecting the baby
- 2. Intrapartum fever A sign that there might be an infection, which raises concern for GBS transmission.
- 3. Rupture of membranes for ≥18 hours The longer the water is broken, the higher the risk of GBS transmission
- 7. Which hormone or inflammatory mediator is elevated in Primary dysmenorrhea?
  - PGF2Alpha
  - In females, PGF2 $\alpha$  is critical for luteolysis, the process of breaking down the corpus luteum, which leads to a drop in progesterone and the start of a new estrous or menstrual cycle.
  - It also induces uterine contractions, so it may be used medically to induce labor or terminate pregnancy.





#### **Psych**

Language 6 months → coos 12 months 1st word

2 year = 50 word vocab, 2 word sentence


3 years = 300 word

Oratory—says "mama" and

"dada" (by 10 mo)

Words—uses 50-200 words (by 2 yr), uses 300+ words (by 3 yr)

Language—understands 1000 (3 zeros) words (by 3 yr), uses complete sentences and prepositions (by 4 yr)



- 1. Oppositional defiant disorder vs conduct?
  - no violence vs violence
  - No violence in ODD and alot of violence in Conduct disorder

- 2. Bupropion is contraindicated in which type of patients?
  - Bulimia and Anorexia as it can cause seizures in these patients --> Pixorize (Shivering = Seizures, Skinny loose pants)
- 3. What is the difference between Social Anxiety disorder (Social Phobia), agoraphobia and schizotypal and panic disorder?
  - Social Anxiety disorder = Fear or embarrassment or rejection ( Like a business meeting or presentation .)
  - Agoraphobia = Intense fear and panic over being in open spaces, crowds or outside homes
  - Schziotypal = prefers to be a loner
  - Panic disorder
- 4. Which ego defence is defined as separating feelings from ideas and events?
  - Isolation of affect (immature defence e.g. describing murder in graphic detail with no emotional response
- 5. What alcohol withdrawal complication occurs between 12 to 48 hours?
  - Withdrawal seizures or Alcohol hallucinations

Alcohol withdrawal syndrome		
Manifestation	Symptoms/signs	Onset since last drink (hours)
Mild withdrawal	Anxiety, insomnia, tremors, diaphoresis, palpitations, gastrointestinal upset, intact orientation	6-24
Seizures	Single or multiple generalized tonic-clonic	12-48
Alcoholic hallucinosis	Visual, auditory, or tactile; intact orientation; stable vital signs	12-48
Delirium tremens	Confusion, agitation, fever, tachycardia, hypertension, diaphoresis	48-96

- 6. How to differentiate between bipolar type 1 and type 2?
  - Bipolar 1
  - --> Manic episodes
  - --> Depressive episodes common but NOT required for diagnosis
  - Bipolar 2
  - --> Hypomanic
  - --> more than or equal to 1 MDD episodes

#### Bipolar & related disorders Hypomanic episode Manic episode · Symptoms less severe · Symptoms more severe ≥4 consecutive days 1 week unless hospitalized · Unequivocal, observable change in · Marked impairment in social or functioning from patient's baseline occupational functioning or · Symptoms not severe enough to hospitalization necessary cause marked impairment or May have psychotic features; makes episode manic by necessitate hospitalization definition · No psychotic features Bipolar I Manic episode(s) · Depressive episodes common, but not required for diagnosis Hypomanic episode(s) ≥1 major depressive episodes required Cyclothymic disorder At least 2 years of fluctuating, mild hypomanic & depressive symptoms that do not meet criteria for hypomanic episodes or major depressive episodes

- 7. What are examples of 4 Hallucinogens?
  - Maraujna, LSD, phencyclidine and MDMA
  - MDMA--> Primarily increases serotonin release (also affects dopamine and norepinephrine).
  - High serotonin levels can lead to altered perception, enhanced sensory input, and emotional openness — features that contribute to its mild hallucinogenic effects.
     PCP--> NMDA receptor antagonist → blocks glutamate signaling in the brain. This causes disruption of normal sensory input, detachment from reality, and profound hallucinations.
- 8. What is the time duration of Brief psychotic disorder?
  - than equal 1 days to less than equal to 30 days More
- 9. What is the time duration of acute stress disorder
  - than equal 3 days to less than equal to 30 days More
- 10. A patient w/ AFIB and delirium can we give me Haloperidol treatment?
  - No we can NOT give haloperidol as it will be detrimental
- 11. Tx of Adjustment disorder ?
  - Psychotherapy
  - Psychotherapy targets maladaptive responses to stress, including the dysfunctional thought patterns contributing to low mood and anxiety (eg, "nothing ever works out," "never going to find anyone") while avoiding unnecessary polypharmacy.
  - Yours = Reassurance

- 12. What is the NBS in patient w/ insomnia in the elderly?
  - CBT → CBT typically combines several approaches, including addressing dysfunctional thoughts about sleep with cognitive therapy, and behavioral measures such as sleep hygiene, sleep restriction, stimulus control, and relaxation techniques.
  - Wrong Answer = Zolpidem at bedtime → The nonbenzodiazepine zolpidem is effective for the short-term treatment of insomnia but is also associated with adverse effects, including cognitive impairment, somnolence, dizziness, and falls. It should not be used as an initial treatment for insomnia in the elderly. When prescribed for elderly patients, the lowest dose should be used.
- 13. What is the first line of acute mania?
  - Antipsychotics

Acute mania	
Clinical features	Elevated, irritable, labile mood     Increased energy & activity, decreased need for sleep     Pressured speech, racing thoughts, distractibility     Grandiosity, risky behavior     Marked impairment, may have psychotic symptoms
Management	Antipsychotics (first- & second-generation)     Lithium (avoid in renal disease)     Valproate (avoid in liver disease)     Combinations in severe mania (eg, antipsychotic plus lithium or valproate)     Adjunctive benzodiazepines for insomnia, agitation

- 14. 2M (1) AMS (2) playing in the garage (3) Multiple episode of emesis (4) tonic clonic seizures What is the cause ?
  - {{c1::Ethylene glycol}} --> encephalopathy, emesis, tachypnea Seizures --> The oxalic acid metabolite binds to serum calcium and forms calcium oxalate; this predisposes to hypocalcemia, which can lead to seizures, tetany, and/or arrhythmias (typical onset ~4-12 hr following ingestion). yours = Organophosphates --> Defecation, Urination, Miosis, Bronchospasm/bradycardia, Emesis, Lacrimation, and Salivation (DUMBELS). Other than emesis, these findings are not present in this patient. Step 3 wrongs
- 15. What is the tx of Serotonin Syndrome?
  - Cyproheptadine blocks H1 receptors, preventing histamine from exerting its effects, which alleviates allergy symptoms such as sneezing, itching, and runny nose.
     Serotonin Receptor Antagonism: It also acts as an antagonist at various serotonin (5-HT) receptor subtypes, particularly 5-HT2A and 5-HT2C receptors
- 16. NBME 7 Block 4

What is the tx of acute mania in a pregnant female?

# {{c1::Quetiapine}}

Wrong answer = Lithium --> crosses the BPB, lithium is associated w/ an increase rate of cardiac malformation

<b>×</b> 1	8.	A 22-year-old female university student is brought to the clinic by her mother 1 day after the patient was dismissed from class because of disruptive behavior. Her professor said she stood up while he was teaching and announced that class was dismissed. The patient has pressured speech and speaks rapidly; she says she feels "wonderful" and has no idea why she is at the clinic. Her mother says the patient has been sleeping only 1 to 2 hours nightly during the past week. The patient says she has been staying up to work on papers and that her family is tracking her movements through her laptop. She says she feels like a puppet controlled by her family and makes accompanying marionette-like gestures with her arms. The mother also says her daughter has become increasingly hypersexual during the past week, and she spent an entire financial aid check purchasing clothing and a new computer. Medical history is significant for asthma treated with an albuterol metered-dose inhaler. The patient has not had any recent alcohol or drug use. She is unsure of the date of her last menstrual period. BMI is 32 kg/m². Vital signs are temperature 36.1°C (97.0°F), pulse 115/min, respirations 20/min, and blood pressure 144/79 mm Hg. Physical examination discloses no abnormalities. Urine pregnancy test is positive. Which of the following is the most appropriate pharmacotherapy?
		A) Clozapine
		B) Lamotrigine
		C) Lithium carbonate
		D) Quetiapine
		E) Valproic acid
		F) No pharmacotherapy; psychotherapy only
C	or	ect Answer: D.
o N o t t	ne isn lar ue ffe ue era era	lar disorder is a mood disorder that features acute episodes of mania. Acute mania typically manifests with an irritable or euphoric mood, increased gy, a decreased need for sleep, increased goal-directed activity, rapid speech and thinking, irresponsibility (as evidenced by this patient's attempt to iss her class), and grandiose thinking. Some patients demonstrate associated psychotic symptoms (eg, grandiose delusions, hallucinations), agement of acute mania requires mood stabilizing therapy (eg, mood stabilizers or mood stabilizing antipsychotics). Atypical antipsychotics (eg, iapine, olanzapine), which antagonize dopamine receptors and modulate serotonin activity, possess mood stabilizing properties and demonstrate stiveness for acute mania over the course of days. Alternatively, for acute bipolar depression, the most effective medication choices are lurasidone, iapine, lamotrigine, and fluoxetine-olanzapine. In patients who are pregnant, careful regimen selection should be undertaken due to the potential ogenic effects of many of the common therapies of bipolar disorder. For this reason, quetiapine is regarded as the treatment of choice because of logenicity. Managing the acute bipolar exacerbation is important as irrational and impulsive behavior during such a state can be harmful to both and fetus.
li	ncc	rrect Answers: A, B, C, E, and F.
		apine (Choice A) is not correct. Clozapine is an antipsychotic used to treat refractory schizophrenia. Its major toxicities include agranulocytosis, whi be severe and life threatening in the setting of infection.

NBME 7 Block 4 11F (1) 1 week of insomnia (2) symptoms began after a bite from a snake (3) snake removed from the house but she still has these symptoms What is the NBS? Correct = {{c1::Reassurance Only}} Yours = Gradual reexposure to snakes while she is in her bedroom --> she has NO fear to snakes, she only has insomnia

<b>x</b> :	34.	An 11-year-old girl is brought to the office by her parents because of a 1-week history of sleeping problems. The symptoms began when her
		brother's snake got out of its cage and crawled into her bedroom and over her foot. She was not bitten or injured and the snake has since been
		removed from the house, but she continues to have difficulty falling asleep and awakens frequently during the night. Her school performance
		continues to be good. Medical history is otherwise unremarkable. She lives with her mother, stepfather, and 8-year-old brother. The patient is 14
		cm (4 ft 9 in; 50th percentile) tall and weighs 28 kg (62 lb; 10th percentile); BMI is 16 kg/m² (25th percentile). Vital signs are temperature
		36.4°C (97.5°F), pulse 76/min, respirations 18/min, and blood pressure 108/68 mm Hg. Physical examination shows no abnormalities. Which of
		the following is the most appropriate management?
		1) 0
		A) Glonazepam therapy

B) Cognitive behavioral therapy

G) Fluexetine therapy

D) Gradual reexposure to snakes while she is in her bedroom

N7B4

Correct Answer: E.

This patient's experience with her brother's pet snake likely resulted in anxiety that is causing sleep difficulty. Stressful events commonly cause temporary emotional problems such as sadness, anxiety, or difficulty sleeping. These problems typically resolve within days to weeks (or in the case of bereavement, months). Adjustment disorder refers to the onset of an emotional disturbance within 3 months of an identifiable stressor. This disorder manifests as marked distress out of proportion to the stressor or a significant impairment in school, work, or social functioning. This patient does not experience dysfunction currently, though her difficulty sleeping even in the snake's absence can be considered out of proportion to the stressor. In any case, adjustment disorder is typically self-limited, so reassurance is appropriate. In severe cases, brief supportive psychotherapy or symptom-focused pharmacotherapy can be beneficial.

Incorrect Answers: A, B, C, and D.

Clonazepam (Choice A) is a benzodiazepine medication commonly used to treat anxiety and rapid-eye movement sleep behavior disorder. Benzodiazepines, though sometimes used, are not first-line agents for acute insomnia. Additionally, the patient does not experience daytime symptoms, and clonazepam is a long-acting medication that would affect the patient during daytime as well.

Cognitive behavioral therapy (Choice B), which commonly includes graded exposure therapy, or gradual reexposure to snakes while she is in her bedroom (Choice D) is used for full-fledged anxiety disorders (eg, specific phobia of snakes). However, this patient does not directly express a fear of snakes or meet duration criteria for specific phobia. Her insomnia is likely to be self-limited and improve without intensive treatment.

7. A 54-year-old man comes to emergency 90 minutes after attending a party in his friend's cellar. He complains of severe stomach cramping and worsening visual disturbance. He appears acutely intexicated. Serum laboratory studies show:

HCO3:: 12 mEq/L Na\*: 142 mEq/L Cl:: 100 mEq/L pH: 7.14

pCO<sub>2</sub>: 23 mm Hg PO<sub>2</sub>: 110 mm Hg

Which of the following is the most appropriate first-line treatment for this patient?

- A) Ethylene glycol
- B) Ethanol
- C) Fenoldopam
- D) Fomepizole
- E) Pralidoxime
- 45. A 29-year-old man with schizophrenia comes to the office for a routine examination. His only medication is risperidone. He does not drink alcohol or use illicit drugs. Vital signs are normal. Physical examination shows no abnormalities. On mental status examination, he has an expansive mood. He pulls out pages of intricately drawn and colored diagrams of triangular shapes and describes the connections among the seven colors of the rainbow, the seven notes on the piano, the seven "special organs of the body," and the epiphanies he has while meditating on the seventh day of the week. He explains that the special organs will resonate when he plays corresponding keys on the piano. He eats foods whose color corresponds to a specific day of the week. As the interview concludes, the patient asks what the physician thinks of his diagrams and theories. Which of the following is the most appropriate physician response?
  - A) "I find them a bit confusing."
  - B) "Tell me more about them."
  - C) "They seem a bit improbable, but I do understand your logic."
  - D) "They seem to represent something important to you."
  - E) "Your ideas are wonderful."

#### Correct Answer: D.

Validation helps build therapeutic alliance. Validation refers to acknowledging a patient's experience as legitimate. In a patient expressing delusional beliefs, as in this case, the physician may not agree with the content of the beliefs. However, expressing disagreement with the content of the beliefs is likely to damage the therapeutic alliance. Conversely, validating the content is disingenuous and may strengthen the delusion. Instead, the physician should strive to identify and validate the patient's experience of the beliefs. Aspects of the patient's experience the physician can validate include the meaning of the experience to the patient or underlying emotions (which the patient may verbally express or indirectly express through body language). In this case, "They seem to represent something important to you" is the most appropriate response from the physician. Building therapeutic alliance is important in promoting treatment adherence and response to treatment.

Incorrect Answers: A, B, C, and E.

"I find them a bit confusing." (Choice A) and "They seem a bit improbable, but I do understand your logic." (Choice C) are overly intellectualized responses that stray from the patient's experience of the belief and are therefore less validating than the correct response. They also challenge the logic of the patient's beliefs, which is likely to damage the therapeutic alliance. Moreover, Choice C moves toward validating the content of the patient's beliefs, which could strengthen his delusions.

"Tell me more about them." (Choice B) avoids answering the patient's question, which could damage the therapeutic alliance. As well, it is clear the patient is experiencing delusional beliefs. Gathering more open-ended information is unlikely to be clinically informative and may decrease time available for treatment planning.

"Your ideas are wonderful." (Choice E) validates the content of the beliefs, which could strengthen his delusions. In addition, this statement strays from the patient's experience of the beliefs and is therefore less validating than the correct response.

Educational Objective: If a patient expresses delusional beliefs, the physician should validate the patient's experience of the beliefs rather than the

#### 32F

- (1) 6 weeks ago she had a sepsis
- (2) today she has fever, chills and weakness
- (3) she is a nurse?

What is the cause of the bacteremia?

- Factitious disorder
- Self-injection of exogenous material
- Yours = Unrecognized IA abscess

UWSA 1 Step 3 Block 1

41F

- (1) insomnia
- (2) Weight loss
- (3) excessive thoughts about death of husband who died 3 weeks ago

dx?

{{c1::B}}rief Grief reaction --> Grief is a common trigger for a MDD. If symptoms intensify into MDD, then SSRI can be given

# Grief vs major depressive episode (MDE)

Grief		MDE
Emptiness from loss	Intense sadness	Hopelessness & misery
Comes in waves (pangs)	Ruminative thoughts	Unrelenting
↓ over days to weeks	Insomnia	Self-loathing, self-hate,
Self-esteem intact	↓ appetite	"I'm worthless"
Suicidal ideation (desire to join the deceased)	Weight loss	Suicidal ideation (desire to end one's pain, "underserving of life")

©UWorld

10 painful bowel movement dialy for 2 weeks

labs are normal

What is the dx?

Factitiotus disorder

Yours = Somatization disorder patient is excessively preoccupied by 1 or more somatic symptoms which is disrupte daily life

Key features of somatic symptom & related disorders		
Somatic symptom disorder	≥1 unexplained symptoms; excessive thoughts, anxiety & behaviors in response to symptoms	
Illness anxiety disorder	Minimal to no symptoms; preoccupation with idea of having a serious illness	
Conversion disorder (functional neurologic symptom disorder)	Neurologic symptom(s) incompatible with anatomy or pathophysiology	
Factitious disorder	Falsification of symptoms/inducing injury in the absence of obvious external rewards	
Malingering	Falsification of illness for obvious external rewards	

# **Inner Circle notes Step 2 and 3 RAPID REVIEW**

# Renal: (1) acid base (2) 5 electrolytes (3) GN (4) Stones (5) Tumors (6) Peds recurrent UTI

- 10. What syndrome is Wilms tumor (nephroblastoma) associated with?
  - WAGR
- 11. What does WAGR stand for ?
  - Willms, Anirdia, Genitourinary abrnomalities and Retardation of the mind
  - Aniridia means lack of an iris.
- 12. For every 1 g decrease of albumin how much does calcium decrease?
  - For every 1 g decrease of albumin calcium decreased by 0.8.
- 13. What is the formula for corrected calcium?
  - Corrected calcium = (measured calcium + 0.8(4 albumin)
  - For each 1 g/dL decrease in albumin, total calcium decreases by ~0.8 mg/dL, even though ionized calcium may remain normal.
- 14. NBEM 6 Block 1 What are the eye abnormalities seen Wilms tumor (nephroblastoma?
  - {{c2::Abnormal pupils}} and {{c1::nystagmus}}

We can get abnormal pupils typically large, irregular or poorly reactive due to the lack of iris musculature. Light sensitivity and poor visual acuity are common. Nystagmus can occur in WAGR syndrome but usually secondary to poor vision from aniridia.

- 1. What is the underlying cause of Pigment Cast?
  - Rhabdomyolysis → myoglobin and other proteins, which can appear pigmented due
    to the heme component of myoglobin. Each myoglobin molecule contains one heme
    group, which allows it to bind one molecule of oxygen.
- 2. When do we give Ca Carbonate in patients w/ hyperkalemia?

- ECG changes and potassium above 6.5 meql/l
- 3. Tx of the acute PSGN?
  - Loop Diuretics → Acute kidney injury (AKI) with decreased glomerular filtration (eg, ↑ serum creatinine) and impaired sodium excretion is also characteristic. The aggregate consequence of both sodium and water retention is volume overload with hypertension and edema. Hence we use loop diuretics for PSGN due to volume overload t
- 4. What type of renal cast are present in a dehydrated person?
  - Hyaline glassy and translucent in appearance → They're made up of a protein call uromodulin and are often the result of conditions such as dehydration and strenuous exercise, as well as chronic kidney failure. Hyaline casts are the most common type of urinary casts, appearing as pale, glassy, and structureless material in urine.




- 6. What does the second A in SAAD of renal papillary necrosis stand for ?
  - Acute Pyelonephritis → Infection-Induced Ischemia: In pyelonephritis, there is an acute inflammatory response. This inflammation can lead to edema and vascular congestion, reducing blood flow to the renal medulla, specifically the renal papillae
- 7. What does the second H in HARDASS mnemonic stand for ?
  - Hypercholermia → hyperchloremic metabolic acidosis results from the loss of bicarbonate (HCO₃⁻), whether through gastrointestinal losses (diarrhea) or renal losses.
- 8. What is the formula for Anion gap Metabolic acidosis?
  - Na ( Cl- + Hco3 -)
  - You though = Na K+ Hco3 -
  - Minimal Contribution: Potassium levels are relatively low compared to sodium and are not considered a major contributor to the overall electroneutrality
- 9. What is the difference between WILMS tumor (nephroblastoma) and neuroblastoma?
  - Does NOT cross the midline WIlms tumor (nephroblastoma) whereas neuroblastoma crosses the midline
- 10. In patients w/ recurrent calcium stones and hypercalciuria what drug helps?
  - {{c1::Thaizide}}
  - Wrong answer = decrease intesintal Ca absorption
- 11. What is a poor long term prognosis in a 40 M w/ PSGN and gross hematuria?
  - {{c1::Age}} Correct = Age of diagnosis Wrong answer = Gross hematuria --> None of these findings significantly affects the prognosis, whereas the difference in prognosis between adults and children has been noted in multiple studie Children: Typically

experience a more benign and self-limited immune response to streptococcal infection. Adults: Often exhibit a more intense and prolonged immune response, which can cause more severe glomerular injury, leading to persistent or progressive renal damage.

# 15. What are the indications for hemodialysis?

Indications for urgent dialysis (AEIOU)		
<u>A</u> cidosis	Metabolic acidosis     pH <7.1 refractory to medical therapy	
<u>E</u> lectrolyte abnormalities	Symptomatic hyperkalemia     ECG changes or ventricular arrhythmias     Severe hyperkalemia     K > 6.5 mEq/L refractory to medical therapy	
Ingestion	Toxic alcohols (methanol, ethylene glycol)     Salicylate     Lithium     Sodium valproate, carbamazepine	
<u>O</u> verload	Volume overload refractory to diuretics	
<u>U</u> remia	Symptomatic:     Encephalopathy     Pericarditis     Bleeding	



# Pulm:

What disease presents the first HAPE or Multifocal pneumonia?

 HAPE due to the acute physiological stress of hypoxia and increased vascular permeability leads to edema formation relatively quickly.

- Why does Multifocal pneumonia occur later? Respiratory infections take time to develop as the immune system weakens due to prolonged hypoxia, dehydration, and cold exposure. Some cases may be due to aspiration or superimposed bacterial infections in people with altitude-related respiratory distress.
- 2. Tx of laryngotracheitis and associated symptoms of SEVERE Croup?
  - nebulized racemic epi ( racemic epinephrine will less likely lead to side effects) stridor at rest Mild to moderate resp distress
- 3. How to dx Vascular Rings in children? rigid bronchoscopy or CT angiogram of the chest
  - CT Angiogram of the chest → Although bronchoscopy would also reveal tracheal indentation in a patient with a vascular ring, - it would not identify abnormal vasculature, making this invasive procedure unnecessary. Diagnostic imaging is the priority when a vascular ring is suspected.
- 4. What are the Ventilator O2 (target) requirements?
  - 88%-95% f-> Preventing hypoxemia and avoid hyperoxia hence o2 toxicity to the organs
- 5. Carcinoid tumors secrete
  - Histamine, serotonin, bradykinin, prostaglandin, substance P and chromogranin
- 6. Cholesterol is high or low in exudates
  - is above 60 mg/dl in exudates due to increase permeability
- 7. Chylothorax: what will be high?
  - Triglycerides --> lymphatic fluid, which normally carries dietary fats, particularly triglycerides, bypasses the normal absorption and drainage pathways and accumulates in the pleural space.
  - Chylos: juice, Chyle --> Chyle is a milky fluid consisting of lymph and emulsified fats, formed in the small intestine during digestion.
- 8. What is another name for Idiopathic Pulmonary Fibrosis (IPF)?
  - Usual interstitial pneumonia
  - Definition: A specific type of chronic, progressive interstitial lung disease of unknown cause.
- 9. What is the 4 parts of the Centor Criteria?
  - CENT = Cervical LAD, Exudates Fever above 38 C no cough
- 10. How long does Chemical Pneumonitis take?
  - Immediately (vs Aspiration Pneumonia which takes longer to develop)
- 11. How to diagnose a patient with suspected Chronic Bronchitis?
  - PFT, Wrong Answer CT scan is a wrong answer for diagnosing chronic bronchitis
  - Purpose of PFT: Assess lung function and airflow obstruction.Reduced FEV<sub>1</sub>/FVC ratio
  - Airflow limitation not fully reversible (hallmark of COPD, which includes chronic bronchitis) Clinical Criteria (very important): Cough with sputum production for at least 3 months in 2 consecutive years, without another identifiable cause.
  - CT Scan of the Chest Supplemental, not first-line
  - Purpose: Evaluate for other lung pathologies (e.g., bronchiectasis, emphysema, lung cancer).
- 12. What is the extubation criteria?

- Adequate oxygenation (eg, PaO 60 mm Hg) on minimal support (ie, fraction of inspired oxygen FiO 40% and positive end-expiratory pressure PEEP 5 cm HO
- Intact inspiratory effort and sufficient mental alertness to protect the airway

What is the intubation criteria?

- Decreased level of consciousness (e.g., GCS ≤ 8), leading to loss of airway protective reflexes
  - 2. Inadequate Oxygenation or Ventilation
- Severe hypoxemia (e.g., SpO<sub>2</sub> < 90% on maximal oxygen)
- Hypercapnia with respiratory acidosis (e.g., pH < 7.25 with high PaCO<sub>2</sub>)
- Signs of respiratory fatigue or failure (e.g., accessory muscle use, tachypnea, low tidal volume
  - The initial criteria for extubation readiness include:



pH >7.25

Adequate oxygenation (eg, PaO ≥60 mm Hg) on minimal support (ie, fraction of inspired oxygen [FiO] ≤40% and positive end-expiratory pressure [PEEP] ≤5 cm HO)

Intact inspiratory effort and sufficient mental alertness to protect the airway

PPPF
pH
PaO2 PaC
PeeP PaC
FiO2 Fi

Ph 7.35 PaO2 94 mmH aCo2 45 mmH FiO2 of 60%

- 13. Noninvasive Positive pressure mechanical ventilation What is the physiological change on the Right and Left heart preload ?
  - RV preload → decrease (increase ITP → decrease CVP return → right heart preload)
  - RV afterload → increase ( pulmonary compression) → Elevated intrathoracic pressure → compresses alveolar capillaries → raising the pulmonary vascular resistance
  - LV preload → PPV decreases MAP through baroreceptor effects and by lowering adrenergic tone through decreasing neurohormonal activation. This is particularly helpful with a superimposed hypertensive emergency, as in this patient.
  - LV afterload → PPV decreases LV transmural pressure because, as the positive pressure is transmitted to the pleura, the LV contracts against a smaller transmural pressure gradient. This is especially beneficial in pulmonary edema, where massive inspiratory efforts generate deep negative pleural pressures to impede LV contraction.
- 14. What is the pathophysiology of BPD?
  - Reduced septation of the alveoli


BPD leads to oxygen toxicity --> fibrotic scarring --> reduced alveoli and disrupted vasculogenesis

Yours = Refeeding induced metabolic disturbance


Step 2 Supplemental

- 15. What do we give for the treatment of exercise induced asthma multiple times a week?
  - Multiple times a week ICS + Formoterol or Antileukotrienes
- 16. What is the tx of exercise induced asthma a few times a week?
  - SABA + antileukotrienes
- 17. How do the cystic fibrosis channel work in the sweat glands versus the lungs?
  - Sweats gland = decrease chloride reabsorption
  - Lungs = decrease secretions of chloride


#### **Sweat ducts**



Respiratory mucosa




- 18. Decrease BS, decrease fremitus and increase resonance on percussion of the apex of left lung. What is the likely diagnosis ?
  - Pneumothorax Or Emphysema




- 19. Polymyositis can develop ILD confirmed on HRCT. What is the NBS in diagnosis?
  - PFT → Pulmonary function tests (PFTs) should be performed to distinguish ILD from respiratory muscle weakness and will show decreased forced vital capacity, total lung capacity, and diffusing capacity.
  - your = Bronchoscopy --> Bronchoscopy with transbronchial biopsy or bronchoalveolar lavage may be needed to diagnose ILD in patients who do not have characteristic findings on CT imaging and PFTs. However, CT and PFTs are usually performed first.

- 20. What are the causes of rightward shift?
  - Increase H+ Increase Temp and increase 2.3 BPG



- 1. Increase temperature = Elevated temperature causes hemoglobin's molecular structure to change slightly, reducing its affinity for oxygen.
- 2. Increase 2,3 DPG = 2,3 DPG binds to haemoglobin and stabilizes the deoxygenated form, making it harder for o2 to bind
- 3. Increase H+ Decreased (pH) =  $H^+$  ions bind to hemoglobin and alter its structure, decreasing its oxygen affinity. This facilitates oxygen release where  $CO_2$  and acid are accumulating (active tissues).
- 1. Decreasing temperature = left shift Cold temperatures stabilize the oxygenated form of hemoglobin. Hemoglobin holds onto oxygen more tightly (higher affinity).
- 2.. Decrease 2,3 DPG --> Leftward shift --> Less 2,3-DPG = less competition with oxygen  $\rightarrow$  hemoglobin binds oxygen more tightly.
- 3. Decrease H+ --> fewer H+less acidic environment = higher pH hemoglobin structure favour the oxygenated state.
- 21. What is the NBS in epiglottitis w/ impending airway compromise?
  - Correct = Intubate the patient, Wrong answer = Epi --> we use neb epi for croup
- 22. How long does it take for fat embolism to occur?
  - Fat Embolism after 24 to 72 hours → Over time, the fat globules are broken down, releasing free fatty acids that cause inflammation and damage to the capillary beds in the lungs (acute respiratory distress syndrome, ARDS) and other organs.
  - The inflammatory response typically peaks around 24–72 hours, leading to the clinical features of fat embolism syndrome.
- 23. What are the histological findings of Asbestosis?
  - Ferruginous bodies are composed of a core fiber (usually asbestos) coated with a golden-brown, iron-containing proteinaceous material.



# <u>cvs</u>

What are other causes of wide pulse pressure?

{{c2::AV malformation }}

{{c1::PDA}}

Wide pulse pressure means blood is quickly leaving the arterial circulation





# Why B is wrong - Bounding pulses

- These means the arterial pulses are *strong*, which is literally the opposite of what we'd expect with an arterial
- ulcer.

   Bounding pulses mean blood is quickly leaving the arterial circulation.

   8/10 Qs on USMLE this is aortic regurg.

   1/10 is PDA. 1/10 is an AV fistula/malformation.







Yours = Medial malleolar ulceration

--> Venous Insufficiency: Chronic venous hypertension can lead to venous stasis ulcers, often located around the medial malleolus due to impaired blood return from the lower limb. Venous Hypertension: In venous insufficiency, faulty or damaged valves in the superficial or deep venous system lead to impaired venous return. This causes blood to pool and increases venous pressure, especially in the lower limbs.

#### Punch out lesion is arterial ulceration

- 1. A patient presents w/ suspected Myocarditis, What is the NBS?
  - Echo
- 2. Cardiac tamponade effect on VR?
  - decrease VR due to increased intrathoracic pressure, Decreased VR > decreased
     CO > increases SVR
- 3. LV MI effect on LV and RV preload -->
  - LV and RV preload increase due to decrease CO more RAAS
- 4. RV MI effect on preload -->
  - RV preload increase due to decrease CO more RAAS
- 5. CVP is related to which heart's preload is right heart or left heart?
  - Right heart
- 6. PCWP is related to which heart's preload is right heart or left heart?
  - Left heart
- 7. What happens to A2 in VSD?
  - Early A2 → L to R Shunting --> increase pulmonary blood flow --> leading to volume overload in the left atrium and ventricles --> increase LV stroke volume --> increase preload leads to accelerated ventricular contractions and emptying. --> early A2
- 8. What happens to CO in Septic Shock? (increase/decrease/ normal)
  - Increase

- Cardiac output is usually elevated.
  - This is due to: (1) **Peripheral vasodilation** → decreased systemic vascular resistance (SVR), (2) **Compensatory tachycardia** and **increased stroke volume**,
  - (3) Activation of catecholamines and inflammatory mediators.
- 9. What happens to CO in Neurogenic Shock? (increase/decrease/ normal)
  - decrease SVR decrease preload
- 10. What happens to SvO2 in Septic shock?
  - Increase because less extraction from tissues
- 11. What is the criteria for a pacing device?
  - 1. LV EF < 35%
  - 2. NYHA Class 2,3,4 HF
  - 3. LBBB w/ QRS duration > 150 msec
  - Normal QRS is 0.06 to 0.10 seconds (60 to 100 milliseconds). A duration longer than 0.10 seconds suggests a delay in ventricular conduction, which can be due to various conditions like bundle branch block or ventricular hypertrophy. In patients with heart failure and LBBB with wide QRS, the heart's ventricles don't contract in a coordinated way. This is called dyssynchrony, and it leads to: Less efficient pumping of blood Worse heart failure symptomsProgressive remodeling and deterioration of heart function
- 12. What are the 2 types of Hypertensive Emergencies?
  - Hypertensive encephalopathy and Malignant Hypertension
  - Malignant Hypertension = Severe HTN w/ retinal hemorrhages, exudates or papilledema
  - Hypertensive Encephalopathy = Severe HTN w/ cerebral edema and non-localizing neurological symptoms and signs
- 13. 52F with a history of shortness of breath, peripheral edema, Jugular venous pressure 9 cm, and crackles both lungs. She had breast cancer 8 years ago for which she takes Doxorubicin and has had radiation therapy. ECHO shows normal EJ fraction. What is the underlying heart defect ? NBME 6 BLock 1
  - A. Endocardial infection w/ spirochete transmitted by an arthropod vector
  - B. Fibrosis of the myocardium causing impaired ventricular filling
  - C. Increased resistance in the pulmonary arteries caused by multiple microthrombi
  - D. Direct cytotoxic effect of chemotherapy drug by on the CMC
  - Correct = Fibrosis of the myocardium causing impaired ventricular filling
  - Yours = Increased resistance in the pulmonary arteries caused by multiple microthrombi
- 14. What type of cardiomyopathy does Doxorubicin cause in the heart?
  - Dilated Cardiomyopathy
- 15. What type of cardiomyopathy does lyme disease cause?
  - Dilated Cardiomyopathy
- 16. What are the 2 sensitive markers for heart failure?
  - Dyspnea and Serum BNP >100 pg/ml
- 17. Where is the murmur heard in coarctation of the aorta?
  - Upper {{c1::left sternal}} border
- 18. Grade 2/6 vibratory systolic murmur best left at the lower left sternal border w/ no radiation Dx?

- **VSD**
- 19. Grade 2/6 harsh systolic murmur best at the left sternal border that radiat to the patient back
  - Coaraction of the aorta

	Diagnostic criteria for anaphylaxis
Ana	phylaxis is likely if there is rapid symptom
ons	et & any 1 of the following criteria:
	Skin/mucosa involvement (eg, hives,
1	lip/tongue swelling) & either hypotension of
	respiratory distress
2	Involvement of ≥2 organ systems after
	exposure to a likely allergen
	Skin/mucosa (eg, hives, lip/tongue
	swelling)
	Respiratory (eg, wheezing, stridor,
	dyspnea)
	Cardiovascular (eg, hypotension,
	tachycardia, syncope)
	Gastrointestinal (eg, abdominal pain,
	vomiting, diarrhea)
3	Hypotension after exposure to a known
3	allergen

- 21. Infective endocarditis ATB prophylaxis
  - Resp iincsion AMox prophylaxis
  - Apical gingiva Amox prophylaxis
  - Infected skin and muscle Vanco Prophylaxis
  - GI and GU procedure in an active infection Prophlyaxis w/ AMp

### **GIT**

### 1.Tx of functional constipation in children?

- Prune juice (prune dried plump w/ high dietary fiber)
- Extra information: Prune juice is a Soluble Fiber: This type of fiber dissolves in water and forms a gel-like substance in the digestive tract. This gel helps to soften the stool, making it easier to pass.
- 2. Tx of Testicular torsion?
  - Orchiopexy → Torsion is the rotation of the testes around the spermatic cord leading to a transversely oriented horizontal testis and high riding testes. Orchidopexy helps restore blood flow to the affected testicle and helps to fix the testis.
- 3. What Is the tx of IBS?
  - Loperamide, hyoscyamine sulfate, Dicyclomine


- Loperamide is an opioid receptor agonist that primarily acts on the μ-opioid receptors in the myenteric plexus of the large intestine. It tx diarrhea
   Hyoscyamine is an anticholinergic (antimuscarinic) agent that blocks the action of acetylcholine at parasympathetic sites in smooth muscle, secretory glands, and the CNS. It reduces gastric secretions, relaxes smooth muscle, and alleviates spasms
- Dicyclomine is an anticholinergic/antispasmodic agent that antagonizes muscarinic receptors in the gastrointestinal tract. It reduces smooth muscle spasms by inhibiting parasympathetic activity, thus relieving abdominal cramping.

#### **Endo**

- 1. What are the 3 key features of Constitutional delay syndrome?
  - (1) delay bone age, (2) family history (3) sexual characteristics delay
- 2. In what trimester is PTU advised to be given ? and In which trimester is MTU advised to be given ?
  - PTU is safe in the first trimester and Methimazole is safe in the 2nd and 3rd trimester
- 3. What is the effect of Dopamine on TSH?
  - Dopamine acts on D2 receptors in the hypothalamus and pituitary. It inhibits the release of thyrotropin-releasing hormone (TRH) from the hypothalamus and directly inhibits TSH release from the pituitary.
- 4. What is the tx of diabetic neuropathy? NBME 6 Block 4
  - Correct = Duloxetine, Wrong answer Amitryptiline --> too many side effects
  - Duloxetine leads to more serotonin and NE in the synaptic cleft which modulates nerve pain signals
- 5. What is the underlying *mechanism* of lithium induced nephrogenic diabetes insipidus?

Acq {{c1::aquaporin}} channel dysfunction

Yours = Vasopressin 2 receptor mutation --> Vasopressin V2 receptor mutations cause inherited AVP-R. However, this condition shows X-linked inheritance; affected male patients can have AVP-R, but female patients are typically asymptomatic.



hypoglycemic episodes due to insulin administration prior to meals  $\check{f}$  ollowed by impaired gastric emptying and delayed absorption

• The indications for **insulin therapy** in diabetes is a newly diagnosed patient with an **A1C >8.5%** or symptomatic diabetes, **pregestational and gestational diabetes**, and **end-stage renal failure**.

Also, if patient has insufficient glycemic control (**target A1c not reached**) over a 3-month period

Also, if patient has insufficient glycemic control (target A1c not reached) over a 3-month period with metform + other antidiabetic drug  $\rightarrow$  initial basal insulin

#### Hematology

# 1. What is the treatment of TACO?

- is Oxygen as well because to much fluid at the alveolar interspace


### 2.Apart from MM and MGUS what other disease is there?

- Waldenstrom MAcroglobulinemia You forgot the name
- 3. What is the normal Mean Corpuscular volume of a RBC?
  - {{c1::80 to 100 Femtoliters }} You always think that 80 to 100 is low MCV
  - "Old people retire at 80-100"
  - Imagine RBCs aging like people.
  - Normal RBC size is between 80 and 100 fL, just like how people commonly retire between 80 and 100 years old

# 4. tx of HUS?

- Blood transfusion and Hemodialysis
- 5. What drug are given for prophylaxis in Sickle cell disease patients?
  - Hydroxyurea encourages the production of fetal hemoglobin, which is less likely to aggregate and form sickle-shaped red blood cells.
  - Vaccination Penicillin and folic acid

6. 85F w/ paleness + 2 months of intermittent dizziness + WBC above 24K, Below is the peripheral smear What is the dx ? NBME 6 Block 3 Q



- Correct = CLL → Numerous lymphocytes w/ large blue nuceli and scant cytoplasm Scant means barely
- Myelodysplastic → BM will show hypercellular marrow w/ abnormal marrow appearing cells of multiple lineage. Periperhal blood smear will show anemai, neutropenia, TCP
- 7. INR monitoring and when to withhold warfarin

Management of supratherapeutic INR			
INR	Bleeding	Treatment	
<4.5	None or minimal	Hold warfarin for 1 or 2 days or decrease dose	Hold Warffarrin
4.5-10	None or minimal	Hold warfarin & resume when INR is therapeutic; administer low-dose (1-2.5 mg) oral vitamin K if there is increased risk of bleeding	Hold Warffarrin Oral Vitamin K
>10	None or minimal	Hold warfarin & administer high-dose (2.5-5 mg) oral vitamin K ORA	Hold Warffarrin L High Does Vitamin K
Any	Serious or life threatening	Hold warfarin & administer intravenous vitamin K 10 mg as well as PCC	Hold Warffarrin V High Does Vitamin K
PCC = prothrombin complex concentrate.			

8. What should be the INR for Warfarin w/ prosthetic valve?

{{c1::2.5 to 3.5 }}

The typical therapeutic INR range for patients on warfarin for most conditions is between 2.0 and 3.0. However, for certain conditions, particularly mechanical heart valves or recurrent thromboembolism, the target INR may be slightly higher, typically around 2.5 to 3.5. This range is decided based on a balance between effectively preventing clot formation and minimizing the risk of bleeding.

9. 2 times Uworld wrong step 3

What is the first change expected to occur in response to tx to IDA?

Correct = increase in {{c1::reticulocyte}} count

Yours = increase in ferrtin levels --> Ferritin reflects iron stores and does not begin to rise until after hemoglobin normalizes.

within 1 to 2 weeks of initiating supplementation the reticulocyte count increases and within a few month the HCT and HG return to normal

Aplastic crisis reticulocytes ?

Less than 1 percent

- Splenic sequestration reticulocytes

More than 1 percent



Cant be C decrease IDA we have LOW ferrtin

What happens to Ferritin and IRON in Anemia of Chronic Disease?

High to normal ferritin cause the body is trying to hide Fe from the pathogen

Low Fe due to increase Hepcidine

• What is the likely *underlying cause* of microcytic anemia in a patient with chronic kidney disease that recently began **erythropoietin therapy**?


#### Iron deficiency

the erythropoietin-induced surge in RBC production can precipitate iron deficiency

#### **MSK**

1. What is the tx of Phantom limb syndrome?

- Multimodal pain regimen like analgesics or opioids or pregabalin (more long term)
- 2. What is the McMurray test used for ?
  - Meniscal tear
- 4. What innervates the teres minor?
  - axillary nerve



- 5. Tx of Sjogren syndrome?
  - Pilocarpine
- 6. What disease has the positive ulnar deviation or Finkelstein's test?
  - quervain tenosynovitis
- 7 45F 6 month hisotry of mild pain and swelling of her right hand

Pain radiates to the shoulder Numbness and tingling in her right thumb and index and ring finergs. Phalen is pos She types alot What is the NBS?

- Correct = Keyboard w/ a wrist pad (splint decrease ROM of the wrist and limits pressure ont eh CT)
- Physical therapy Not decrease compression
- 8. What is the difference between rotator cuff tear and rotator tendinopathy?
  - rotator cuff tear ( weakness above 40) vs rotator tendinopathy (pain)
- 9. What are the features of complex regional pain syndrome?
  - Recent {{c2::trauma}}
  - {{c1::Vasomotor}} changes
  - Severe Paina and {{c3::burning}}/stinging
  - Trophic skin changes
- 10. How is the dx spinal cord compression?
  - {{c1::MRI }}
  - Yours = 99m Technetium bone --> helps in osteomyeltis.
- 11. When do we use 99 m technetrium bone scan?
  - Osteomyelitis
  - <sup>99</sup>mTc-labeled diphosphonates (e.g., methylene diphosphonate, or MDP) are injected intravenously.

These compounds have a high affinity for **hydroxyapatite** in bone—especially in areas with **active bone turnover**.

In osteomyelitis, the infected area has:

- Increased blood flow
- Increased osteoblastic activity
- Bone remodeling
- 13. What is the initial tx of spinal cord compression?
  - Correct = {{c1::High dose dexamethasone}}--> helpful as a bridge to definitive therapy can help control pain and may transiently improve neurologic function in paitnes secondary to decreasing edema at the siter of spinal cord compression.
  - Yours = Radiation therapy --> this is definitive tx of neoplastic epidural spinal cord compression
  - NBME6 Block 3
- 14. What is the NBS in a patient w/ CTS?
  - Serum TSH, Yours RF → rheumatoid factor causes my joint problems but hypothyroidism causes weight gain and accumulation of mucopolysaccharides which leads to CTS ont he NBME 6
- 15. What are the nbs in a patient w/ Spondylolysis?
  - Activity modification
  - Patients with confirmed spondylolysis should limit activity (ie, abstain from sports) and continue conservative management (eg, analgesics, ice/heat) for 90 days to allow the fracture to heal. After 90 days, patients who improve can gradually go back to their activity.
  - Yours = Sruger those who do NOT improve then MRI and spine surgery are advised.

Spondylolisthesis			
Pathophysiology	Bilateral pars interarticularis defects (eg, fractures) → anterior slippage of vertebral body     Most common: L5 slips over S1		
Risk factors	Repetitive back extension & rotation (eg, gymnasts, divers)     Adolescent growth spurt		
Clinical features	Low back pain that is worsened by extension     Radiculopathy as slippage progresses     Palpable step-off may be present		
Diagnosis	Lumbar x-rays (typically visible on lateral views)		
Management	Activity modification (avoid inciting sports), pain control     Neurologic deficits or symptoms >90 days: obtain MRI of spine & surgical consultation		

Apart from Mcmarry test what other test can be used to diagnose meniscus teear?

### Thessaly test for meniscal injury



Patient rotates body side to side with knee in 20° flexion & foot planted

- \_ ©UWorld
- The Thessaly test is a clinical examination used to help diagnose meniscal tears in the knee. It involves rotating the knee while the leg is slightly bent, and a positive test is indicated by pain or a feeling of locking or catching in the joint line.
- 16. What is main risk factor osteoporosis in a patient a 73 M who drinks 1 glass alcohol per day
  - {{c1::Age}} Age: Risk increases with age, especially after menopause in women.
  - Yours = Alcohol intake --> g, ≥3 drinks/day), likely due to decreased osteoblast activity. This
    patient's light alcohol consumption is not likely a significant contributor to osteoporosis
    riskStep 3 wrong
  - As men age, their bone mineral density gradually decreases, especially after the age of 70. This decline is due to reduced osteoblastic activity (bone formation) and increased osteoclastic activity (bone resorption).
- 17. What is the NBS in a patient w/ a dirty wound + who is immunized?
  - {{c1::Nothing}} Yours = He does not require tetanus prophylaxis because his wounds have been cleaned thoroughly
- 18. What is the tx of scleroderma?
  - {{c2::captopril}} and IV {{c1::nitro}} Scleroderma involves widespread microvascular damage and endothelial cell dysfunction. This injury leads to impaired regulation of blood flow and vascular tone. ACE inhibitors are the agents of choice in the treatment of scleroderma renal crisis because they reverse angiotensin-induced vasoconstriction.
- 19. What is the tx of Polymyalgia Rheumatica w/o no symptoms of GCA?
  - {{c2::Low}} dose {{c1::Oral glucocoritcoids }}
- 20. What age is genu varum?
  - 0 to 2 years
- 21. What age is genu valgum?
  - 2 to 6 years
- 22. What is a red flag for scliosis?

- {{c1::Back pain }} causing notrunal awakening --> back pain that causes nocturnal awakening is concerning for an underlying spinal cord tumor. A spinal cord tumor can extend and disturb surrounding nerves, causing constant pain that is described as "dull" or "gnawing." In addition, impingement of spinal cord pathways can cause dysesthesias, weakness, and neurogenic bladder/bowel.
  - Back pain
  - Neurologic symptoms
    - Rapidly progressing curvatur
  - >10 degrees each year) Vertebral anomalies on x-ray
  - Night pain unrelieved by rest can be a sign of bone lesions, metastasis, infection (osteomyelitis, discitis), or malignancy.

#### 23. What are the 5 features of Compartment syndrome?

#### 1. Paresthesias

- 2. Pain out of proportion/on passive stretch
  - 3. Pulse deficit
    - 4. Pallor
    - 5. Paralysis

#### Why Pulse Deficit is Late

- The pressure has to become very high (exceeding arterial pressure) to compress larger arteries and result in a pulse deficit.
- By this stage, muscle and nerve tissues are already suffering irreversible damage (usually after ~6 hours of ischemia).
- Therefore, waiting for a pulse deficit before diagnosing or treating compartment syndrome results in delayed intervention and poor outcomes, including possible limb loss.
- Paresthesia → The increased pressure compresses the neurovascular structures, particularly the nerves running through the affected compartment.

#### 24. How to measure compartment pressure?

- Delta Pressure = Diastolic Blood Pressure Intra-Compartmental Pressure
- Why diastolic BP?
- Perfusion pressure in tissues depends on the pressure gradient between arterial
  inflow and venous outflow, which primarily occurs during diastole when the arteries
  are less compressed and blood flow is maintained. Systolic BP is the peak pressure
  during ventricular contraction and is less relevant for tissue perfusion during the
  diastolic phase.

# 25. WHat does the compression to the LFC nerve lead to?

#### Meros pain thigh, algia = pain



#### **Derm**

- 1. What is Nevus Flammeus?
  - Respect midline (formal) Does NOT gress
- 2. Nevus Simplex
  - Goes away
- 3. What type of Erythema in Acute Rheumatic fever?
  - Marginatum → Marginatum It describes the characteristic appearance of the skin lesion, which has a distinct, well-defined border or margin.
- 4. Which of the following factors in a patient's history or PE is most indicative for further evaluation in an infant w/ infantile hemangiomas?
  - >5 cutaneous lesions --> Liver US
  - Facial/ segmental --> ECHO and MRI of the head
  - Cervical facial --> laryngoscopy
  - Lumbosacral --> spinal US
  - Infants with multiple cutaneous hemangiomas have a statistically higher chance of also having hemangiomas in internal organs, particularly the liver.
- 5. What are desmoid tumors?
  - Desmoid tumors are slow-growing and locally aggressive benign neoplasms with a high rate of local recurrence, even after surgical excision.
- 6. What is the first line tx of atopic dermatitis ?NBME 6 B4
  - PetroPetrolatum based moisturizing ointments
- 7. What is the 2nd line tx of atopic dermatitis?
  - Tacrolimus NBME 6 B4
- 8. What is the tx of Pemphigoid gestationis?
  - topical triamcinolone (triamcinolone is a synthetic corticosteroid, Binding to glucocorticoid receptors: Triamcinolone binds to specific receptors in the cell, called glucocorticoid receptors (GR).)

- 9. Tx of actinic keratosis?
  - 5FU and cryotherapy
- 10. What is the NBS in bullous pemphigoid?
  - Skin biopsy --> Performing a skin biopsy, including both routine histology and direct immunofluorescence, helps confirm the diagnosis of bullous pemphigoid by demonstrating the characteristic subepidermal blisters and immune deposits along the basement membrane, thereby differentiating it from other blistering disorders.

11.



11. A 58-year-old man comes to the office because of a 1-year history of a slowly enlarging, painless, non-itchy lesion on his right inner thigh. He notices the lesion only when he bathes and believes he might be developing a similar one on his right forearm. He has HIV infection well controlled with antiretroviral therapy. He also takes a daily multivitamin. He does not smoke cigarettes or drink alcohol. He used illicit intravenous heroin for 5 years but stopped 35 years ago. He has worked on a farm for 40 years. Vital signs are within normal limits. A photograph of the lesion over the right thigh is shown. Examination of the right forearm shows a 3-cm, raised lesion with well-demarcated borders and dark pigmentation. The remainder of the examination shows no abnormalities. Which of the following historical factors most likely increased this patient's risk for developing these lesions?





49. A 71-year-old man comes to the office with his wife for an initial visit because of a non-itchy, painless spot on the patient's back that the patient thinks has been present for several years. His wife notes that the spot has enlarged and become darker during the past several months. The patient's medical history is unremarkable and he takes no medications. Vital signs are normal. Physical examination discloses no adenopathy. Examination of the patient's back discloses the lesion shown in the photograph. The remainder of the physical examination discloses no abnormalities. Which of the following is the most appropriate next step in management?


- A) Biopsy of the lesion
- B) Cryotherapy
- C) Microscopic examination of a scraping of the lesion
- D) Surgical excision of the lesion
- E) Topical triamcinolone therapy
- F) Observation only

Correct Answer: F

### **CNS**

- 1. tx of Idiopathic Intracranial Hypertension?
  - Acetazolamide due to decrease CSF and topi (increase GABA)
- 2. Dx of Carotid Artery Dissection?
  - W/ CT angiography we can see extra and intra cranial and NOT duplex US because duplex US will not allow us to see the intramural hematoma, collaterals or other stuff
- 3. What happens to Pregnancy associated plasma protein A in Down Syndrome?
  - {{c1::Decreased}}

- Placenta's ability to produce hormones and proteins like PAPP-A could be impaired in pregnancies affected by Down syndrome.
- 4. In Alzheimer disease which lobe do we atrophy of?
  - Temporal lobe
  - We have 2 two hippocampi, one located in each hemisphere of the brain. Each
    hippocampus is situated in the medial temporal lobe. The temporal lobe, particularly
    the hippocampus, is highly active and involved in memory formation and learning.
  - Amyloid-beta plaques and neurofibrillary tangles tend to accumulate initially in the
    entorhinal cortex and hippocampus. Tau proteins are microtubule-associated proteins
    primarily found in neurons, where they help stabilize microtubules involved in
    maintaining cell structure and facilitating intracellular transport.
- 5. TOADG --> what does A stand for ?
  - Vitamin A, Vitamin A increase CSF production
- 6. What type of Nystagmus do we NOT see in peripheral vertigo?
  - Vertical nystagmus is never seen ( due to the orientation of the semicircular canal)
- 7. How long does it take for there to be **dementia** in **Parkinson disease** after motor symptoms?
  - Years → Parkinson's disease primarily affects movement due to the degeneration of dopamine-producing neurons in the substantia nigra. While cognitive decline can occur, it usually involves different areas of the brain, particularly the cortex and subcortical regions. This process of degeneration can take time, often spanning years, before cognitive symptoms become evident.
- 8. What is the management of IIH? Do we have LP or Acetazolamide first?
  - LP before Acetazolamide in IIH



- 9. What does A and B represent?
  - Pyramids which carry the corticospinal tracts?
- 10. What does C and D carry?
  - medial lemniscus Which is for the loss of fine touch, vibration, and proprioception
- 11. What is the difference between Fasciculus Cuneatus vs Fasciculus gracilis?


- Fasciculus Cuneatus = sensory information from the upper limbs, including the arms, upper trunk, and neck
- Fasciculus Gracilis = Carries sensory information from the lower limbs

#### 12. Fever + right tympanic membrane is NOT visible due to copious purulent drainage

What is the dx?

Correct = {{c1::Suppurative otitis media }}

Yours = Serous otitis media occurs when an effusion is present in the middle ear canal in the absence of infection and inflammation. It typically follows episodes of suppurative otitis media, but it is not associated with fever, pain, crankiness, tympanic membrane rupture, or purulent drainage.



- 13. Step 3 wrongs What is the long term guidance for a patient w/ Seborrheic dermatitis?
  - Chronic {{c1::relapsing}} condition
  - Your = 1 course of tx usually leads to long term remission

initial treatment can provide significant improvement in symptoms, but patients usually benefit from intermittent re-treatment. As a result, patients usually benefit from intermittent re-treatment. Typical regimens include topical ketoconazole or ciclopirox every 1-2 weeks.


- 14. What is the NBS in a patient w/ Corneal laceration? (Step 3 wrongs)
  - Correct = consultation w/ {{c1::opthal}}

Yours = Irregation of the eye w/ saline --> Saline irrigation is commonly used to clear a superficial foreign body. In patients with full-thickness corneal lacerations, as in this patient, saline irrigation is contraindicated because it can push environmental pathogens from the surface of the eye into the globe, increasing risk of infection.

Corneal laceration can cause an open globe injury if it extends through the full thickness of the cornea. A positive Seidel sign (ie, concentrated fluorescein uptake with subsequent clearing in a waterfall pattern) indicates full-thickness corneal laceration, an ophthalmologic emergency that requires immediate ophthalmologic consultation.

- 15. What is the NBS in patients w/ keratitis ?Step 3 wrong
  - {{c1::Opathlamologist referral}} --> irsk of blindness hence referral is important

 Yours = Switch to oral fluoroquinolone --> Intravenous or oral fluoroquinolone therapy does not provide any additional drug delivery to the cornea or conjunctiva and does not provide sufficient coverage against gram-positive organisms. However, intravenous therapy may be indicated for patients whose compliance with oral or topical therapy cannot be assured



Plagio = Slanted and cephalic = head

19. Cause of LBBB?


IHD- Myocardial infarction (especially affecting the interventricular septum) can damage the conduction pathways, leading to LBBB, Wrong = LVHT

#### **Short subjects**

- 1. When do we use praziquantel for tx?
  - Schistosomiasis and tinea solium and Diphyllobatrum latat
- 2. Duchenne muscular dystrophy at what age?
  - Below age of 5
- 3. Hepatitis B causes which disease?
  - Hepatocellular cancer
- 4. Most common immunodef is?
  - IgA deficient
- 5. Who will have a positive PPD of above 10 mm will be positive? (4)
  - Health care workers, prison workers, high risk TB reactivation ( DM and Cortico)
- 6. Tetanus prevents the release of which NT?
  - Glycine and GABA
- 7. MArfan syndrome he is asymptomatic. What is the nba prior to sports participation?
  - Echo —> Yours = Slit lamp exam —> not for sports
- 8. 25M w/ 2 weeks of 3rd degree burns and has been on orotracheal ventilation for 10 to 14 days. What is the NBS?
  - Tracheostomy —> improve WOB, earlier ventilatory weaning and improve communication and decrease sedation needs decrease risk of laryngeal injury increase comfort and mobility
- 9. Which cells is Hyperacute rejection mediated by ?
  - Preformed antibodies
- 10. What is the treatment of Hyperacute rejection?
  - Graft removal

#### 11. What is the tx of Invasive Candida or Invasive Aspergillosis

- Echinocandins
- **12.** What is used for the diagnosis of Candida vulvovaginitis?
  - KOH preparation (wet mount is used for a variety of condition but is NOT that sensitive
- 13. MOA of Echinocandins?
  - Inhibition of  $(1\rightarrow 3)$ -β-D-glucan synthase, an essential enzyme involved in the synthesis of  $(1\rightarrow 3)$ -β-D-glucan, a major component of the fungal cell wall.
  - Disrupts fungal cell wall integrity, leading to osmotic instability and ultimately cell lysis and death.
- 14. What are the last 2 steps of fungal cell wall synthesis?



- 15. What does Amphotercin bind to?
  - It binds to ergosterol
- 16. What drug inhibits squalene epoxidase?
  - Turbenfine

## 17. What is the prophylaxis criteria for lyme disease?

TIP BN must meet all 5

- Tick attacted for > 36 hours
- Ixodes scapularis (deer tick)
- Prophylaxis started with 72 hours of tick removal
- Borrelia burgerdogeri infection rate > 20 % for example patient located in New ENgland
- No CI to Dox
- 18. 6 month old infant w/ bronchiolitis what is the infant at greatest risk of developing?


- Correct = {{c1::Recurrent wheezing }}
- Yours = FTT --> dehydration and WL during the symptomatic period due to insensible fluid loss (eg tachypnea) or decreased fluid intake. However, after the infection is resolved the weight gain and growth are expected.
- Infants age <2 months and those born prematurely or with cardiopulmonary disease are at greatest risk for apnea and/or respiratory failure during peak symptoms. In contrast, older, full-term, healthy infants and young toddlers typically recover with no acute complications.



After a burn?

Pseudo

HY USMLE Q #974 - Micro / Derm HY USMLE Q #962 - Micro / Pharm



Sliver stain w/ yeast, methadone  $\rightarrow$  IVDU  $\rightarrow$  HIV  $\rightarrow$  PCP


Lysosomal storage	Lysosomal enzyme deficiency → accumulation of abnormal metabolic products. † incidence of
diseases	Tay-Sachs. Niemann-Pick, and some forms of Gaucher disease in Ashkenazi Iews.

diseases	Tay-Sachs, Niemann-Pick, and some to	ornis of Gaucher disease if	i Asiikeiiazi jews.		
DISEASE	FINDINGS	DEFICIENT ENZYME	ACCUMULATED SUBSTRATE	INHERITANCE	
Sphingolipidoses					
Progressive neurodegeneration, developmental delay, hyperreflexia, hyperacusis, "cherry-red" spot on macula [A. (lipid accumulation in ganglion cell layer), lysosomes with onion skin, no hepatosplenomegaly (vs Niemann-Pick).		Hexosaminidase A GM2 ganglioside. AR     ("TAy-Sax").  Anglokeratomas are hard bumps on your skin. The condition occurs when tiny blood vessels called capillaries get bigger or break open near your skin's surface and the surface of your skin thickens. Red, blue, purple or black bumps form undermeath your skin, Anglokeratomas are a type of noncaneurous (belengt) tumor.			
Fabry disease	Early: triad of episodic peripheral neuropathy, angiokeratomas [3], hypohidrosis.  Late: progressive renal failure, cardiovascular disease.	② α-galactosidase A.	Ceramide trihexoside (globotriaosylce- ramide).	XR	
Metachromatic leukodystrophy	Central and peripheral demyelination with ataxia, dementia.	3 Arylsulfatase A.	Cerebroside sulfate.	AR	
Krabbe disease	Peripheral neuropathy, destruction of oligodendrocytes, developmental delay, optic atrophy, globoid cells.	Galactocerebrosi- dase (galactosylce- ramidase).	Galactocerebroside, psychosine.	AR	
Gaucher disease	Most common. Hepatosplenomegaly, pancytopenia, osteoporosis, avascular necrosis of femur, bone crises, Gaucher cells [ (lipid-laden macrophages resembling crumpled tissue paper).	• Glucocerebrosidase (β-glucosidase); treat with recombinant glucocerebrosidase.	Glucocerebroside.	AR	
Niemann-Pick disease D	Progressive neurodegeneration, hepatosplenomegaly, foam cells (lipid-laden macrophages) □, "cherry-red" spot on macula □.	<b>⊙</b> Sphingomyelinase.	Sphingomyelin.	AR	
Mucopolysaccharidoses					
Hur <mark>l</mark> er syndrome	Developmental delay, skeletal abnormalities, airway obstruction, corneal clouding, hepatosplenomegaly.	α-L-iduronidase.	Heparan sulfate, dermatan sulfate.	AR	
Hunter syndrome	Mild Hurler + aggressive behavior, no corneal clouding.	Iduronate-2 (two)- sulfatase.	Heparan sulfate, dermatan sulfate.	XR	
Sulfatides	GM, Ceramide trihexoside  O  GM, O  Glucocerebroside  O  Ceramide  Sphingomyelin ₪		clearly (no corneal clou aim for the X (X-linke		

ГҮРЕ	INHERITANCE	PATHOGENESIS	† BLOOD LEVEL	CLINICAL
l—Hyper- chylomicronemia ye	AR ea tum nai ghala	Lipoprotein lipase or apo CII deficiency	Chylomicrons, TG, cholesterol	Pancreatitis, hepatosplenomegaly, and eruptive/pruritic xanthomas (no † risk for atherosclerosis). Creamy layer in supernatant supernatant = usually clear
cha infla are	AD Absent or defective IIa: LDL, cholesterol IIb: LDL, cholesterol, defective apo B <sub>100</sub> VLDL  Absent or defective IIb: LDL, cholesterol, defective apo B <sub>100</sub> VLDL  Absent or defective apo B <sub>100</sub> IIb: LDL, cholesterol, vLDL  Absent or defective apo B <sub>100</sub> VLDL  Absent or defective IIa: LDL, cholesterol, vLDL  Absent or defective Absent IIIa: LDL, cholesterol, vLDL  Absent or defective Absent IIIa: LDL, cholesterol, vLDL  Absent		Heterozygotes (1:500) have cholesterol ≈ 300 mg/dL; homozygotes (very rare) have cholesterol ≥ 700 mg/dL.  Accelerated atherosclerosis (may have MI before age 20), tendon (Achilles) xanthomas, and corneal arcus.	
III—Dysbeta- lipoproteinemia APO	AR B NBME g	ApoE (defective in type thrEE)	Chylomicrons, VLDL	Premature atherosclerosis, tuberoeruptive and palmar xanthomas.
IV—Hyper- triglyceridemia	AD	Hepatic overproduction of VLDL	VLDL, TG	Hypertriglyceridemia (> 1000 mg/dL) can cause acute pancreatitis. Related to insulin resistance.

19. Yea tum nai ghalat keeya

I-cell disease (inclusion cell disease/mucolipidosis type II)—inherited lysosomal storage disorder (autosomal recessive); defect in N-acetylglucosaminyl-l-phosphotransferase → failure of the Golgi to phosphorylate mannose residues (↓ mannose-6-phosphate) on glycoproteins → enzymes secreted extracellularly rather than delivered to lysosomes → lysosomes deficient in digestive enzymes → buildup of cellular debris in lysosomes (inclusion bodies). Results in coarse facial features, gingival hyperplasia, corneal clouding, restricted joint movements, claw hand deformities, kyphoscoliosis, and ↑ plasma levels of lysosomal enzymes. Symptoms similar to but more severe than Hurler syndrome. Often fatal in childhood.



Signal recognition particle (SRP)—abundant, cytosolic ribonucleoprotein that traffics polypeptide-ribosome complex from the cytosol to the RER. Absent or dysfunctional SRP → accumulation of protein in cytosol.

#### Vesicular trafficking proteins

- COPI: Golgi → Golgi (retrograde); cis-Golgi
   → ER.
- COPII: ER → cis-Golgi (anterograde). "Two (COPII) steps forward (anterograde); one (COPI) step back (retrograde)."
- Clathrin: trans-Golgi → lysosomes; plasma membrane → endosomes (receptor-mediated endocytosis [eg. LDL receptor activity]

20. What is the prognosis of pertussis?

Correct = {{c1::good prognosis He likely recover completely w/ appropriate tx }}

Yours = He will have life time immunity to this conditon after it resolve --> Generally protection from pertusis, whether in the form of vaccination or naturally acquired immunity wanes over time. For this reason, the booster are recommended every 10 years.

Pertusis occurs as a result of an infection from Bordetella pertussis. Pertussis may lead to severe paroxysmal coughing which may be accompanied by cyanosis and followed by syncope, emesis or apnea

Memory B cells and T cells specific to B. pertussis decline over time, reducing the ability to mount a strong response upon re-exposure.

#### 21. Malaria prophylaxis

Recommended antimalaria chemoprophylaxis options for short-term travelers					
	Medication	Adverse effects/comments			
	Atovaquone- proguanil	Expensive     GI disturbance (eg, abdominal pain), † liver function tests			
Areas with chloroquine- resistant P falciparum (Sub-Saharan Africa, southern & Southeast Asia)	Doxycycline	Inexpensive     Gl disturbance, sun sensitivity, teratogenic			
	Mefloquine	Neuropsychiatric effects     Agent of choice in pregnancy     Weekly dosing			
Areas with <b>chloroquine-</b> <b>susceptible</b> <i>P</i> <b>falciparum</b> (in addition to options above)	Chloroquine, hydroxychloroquine	Need to be started 1-2 weeks in advance     Potential exacerbation of some skin conditions     Weekly dosing			
Areas without <i>P falciparum</i> (parts of South America, Mexico, Korean peninsula)	Primaquine	Potential teratogenicity     Hemolysis in patients with G6-PD deficiency     Weekly dosing			

GI = gastrointestinal; G6-PD = glucose-6-phosphate dehydrogenase.

#### 22. How to prevent Giardiasis?

<sup>©</sup>UWorld

Correct = {{c1::S}}symptomatic students w/ positive stool samples should refrains from using recreational water venus

Transmission occurs via fecal oral route or ingestion of contaminated food/water. Exposure to untreated water during camping or hiking is a common source of infection.

Symptoms of acute infection include bulky, greasy stools.

Recreational water venues are another common source of outbreaks, as cysts often resist chlorination and remain viable in water. Affected individuals should therefore refrain from attending these venues until symptoms have resolved for 2 weeks.

Chronic infection leads to

Yours = Hand sanitizer

Hand hygiene is important in controlling the spread of giardiasis. Soap-and-water-based agents are preferred, as Giardia cysts are resistant to many alcohol-based hand sanitizers and disinfectants.

10M (1) myalgias and arhalgias (2) 38.8 C (3) Petechia on the bilateral wrists and ankles (4) Low platelet (5) camping trip in North Carolina

What is the diagnosis?

Correct = {{c1::RMSF}}

Yours = No therapy is indicated --> Mono

**CLaSH** 

Carcinomas originate from epithelial cells lining organs. Carcinoma cells tend to invade nearby lymphatic vessels because the lymphatic system provides a network of channels that facilitate tumor cell dissemination. Lymphatic vessels are often abundant near epithelial tissues.

Sarcoma originate from mesenchymal tissues such as bone, muscle, cartilage, or connective tissue. Sarcoma cells invade blood vessels directly because of their proximity and the rich vascular supply of mesenchymal tissues.

SKin manifestation of hep C

LCCP = Lichen planus, Cryoglobulinemia, Cutaneous Leuko, PCT

What are other catalse positive organism apart from SPAC?

Serrteria and Bukholdreia

Prophylaxis of CGD

Itraconazole inhibits fungal ergosterol synthesis, inhibits Aspergillus

# TMP-SMX inhibits bacterial folate synthesis, providing broad-spectrum bacterial prophylaxis Prophylaxis against bacterial infections, particularly Staphylococcus aureus

INF gamma: Immunomodulatory therapy to enhance phagocyte function.IFN-γ stimulates the oxidative burst activity of phagocytes, partially compensating for the defective NADPH oxidase.


25. 25M

(1) bite by a dog

What is the NBS?

{{c1::observe dog for 10 days }}

Wrong answer = Sarcific dogs



General Pediatrics, Geriatrics and Public health

- 1. What is considered a High Blood pressure in a 5 year old child?
  - 95th percentile (roughly around 110 mmHg systolic is considered high in children)
- 2. What happens to the renal system in the elderly?
  - reduction in functional {{c2::glomeruli}} Decreased renal {{c1::blood flow }}Limited hormonal responsiveness (eg renin PTH )

- 3. Which age is the HPV vaccine given?
  - 9 to 26
  - You though = 11 to 26 years
- 4. PPSv23 vaccine involves b or t cells?
  - Only b cells, T cell independent → Polysaccharide antigens directly stimulate B cells that have B-cell receptors specific for the polysaccharide.

# This activation bypasses T-cell help, leading to:

- Production of mainly IgM antibodies
- Limited formation of memory B cells
- 5. What are the 2 dosages for Meningococcal vaccine?
  - First dose 11/12 years
  - Second dose: 16 y/o